

# Birkhoff-Kellogg type results with applications

Gennaro Infante

We present some classical and recent results of Birkhoff-Kellogg type in cones. We illustrate their applicability in the context of ordinary, functional and partial differential equations subject to local, nonlocal and functional boundary conditions.

## References

- [1] G. D. Birkhoff and O. D. Kellogg, Invariant points in function space, *Trans. Amer. Math. Soc.*, **23** (1922), 96–115.
- [2] A. Calamai and G. Infante, An affine Birkhoff–Kellogg type result in cones with applications to functional differential equations, *Mathematical Methods in the Applied Sciences*, **46** (2023), 11897–11905.
- [3] A. Calamai and G. Infante, On fourth order retarded equations with functional boundary conditions: a unified approach, *Discrete and Continuous Dynamical Systems - Series S*, doi: 10.3934/dcdss.2023006.
- [4] A. Calamai and G. Infante, Nontrivial solutions of a parameter-dependent heat flow problem with deviated arguments, to appear in: “Topological Methods for Delay and Ordinary Differential Equations”, P. Amster and P. Benevieri (Eds.).
- [5] A. Calamai, G. Infante and Jorge Rodríguez-López, A Birkhoff–Kellogg type theorem for discontinuous operators with applications, submitted, arXiv:2401.16050.
- [6] G. Infante, Eigenvalues of elliptic functional differential systems via a Birkhoff-Kellogg type theorem, *Mathematics*, **9** (2021), n. 4.
- [7] G. Infante, On the solvability of a parameter-dependent cantilever-type BVP, *Appl. Math. Lett.*, **132** (2022), 108090.
- [8] M. A. Krasnosel’skiĭ and L. A. Ladyženskii, The structure of the spectrum of positive nonhomogeneous operators, *Trudy Moskov. Mat. Obšč.*, **3** (1954), 321–346.

**First Author:** Gennaro, Infante

**Affiliation:** *Dipartimento di Matematica e Informatica, Università della Calabria  
87036 Arcavacata di Rende, Cosenza, Italy*

**e-mail:** gennaro.infante@unical.it